

UECM1304 TEST 2 MARKING GUIDE

Name: _____ Student ID: _____ Mark: _____ /20

COURSE CODE & COURSE TITLE: UECM1304 DISCRETE MATHEMATICS WITH APPLICATIONS

FACULTY: LKC FES, UTAR COURSE: AM

TRIMESTER: SAMPLE LECTURER: LIEW HOW HUI

Instruction: Answer all questions in the space provided. **If you do not write your answer in the space provided, you will get ZERO mark.** An answer without working steps may also receive ZERO mark.

1. CO3: Demonstrate various proof-techniques.C3

(a) Use direct proof to show that there are integers m and n such that $7684m + 15283n = 17$.

[**Note:** You need to use extended Euclidean algorithm to find out m and n rather than simply guessing. Guessing the answer is regarded as cheating and only 0.5 marks will be awarded.] (4 marks)

Ans.

$$15283 = 7684 \times 1 + 7599$$

$$7684 = 7599 \times 1 + 85$$

$$7599 = 85 \times 89 + 34$$

$$85 = 34 \times 2 + 17$$

$$34 = 17 \times 2 + 0$$

[1.5 marks]

$$17 = 85 - 34 \times 2$$

$$17 = 85 - (7599 - 85 \times 89) \times 2 = 85 \times 179 - 7599 \times 2$$

$$17 = (7684 - 7599) \times 179 - 7599 \times 2 = 7684 \times 179 - 7599 \times 181$$

$$17 = 7684 \times 179 - (15283 - 7684) \times 181 = 7684 \times 360 - 15283 \times 181$$
 [2 marks]

Therefore, there is $m = 360$ and $n = -181$ such that $7684m + 15283n = 17$ [0.5 mark]

- (b) Let A and B be two sets. Use contrapositive proof to show that if $A \times B = \emptyset$, then $A = \emptyset$ or $B = \emptyset$. (2 marks)

Ans. Suppose $A \neq \emptyset$ and $B \neq \emptyset$ [0.5 mark]

There exist $x \in A$ and $y \in B$ [0.5 mark]

Therefore $(x, y) \in A \times B$. So $A \times B \neq \emptyset$ by definition. [1 mark]

- (c) Use the Principle of Mathematical Induction to prove the equality

$$2^3 + 3^3 + \cdots + n^3 = \left[\frac{n(n+1)}{2} \right]^2 - 1.$$

for integers $n \geq 2$. (3 marks)

Ans. Base Step:

When $n = 2$, RHS = $\left[\frac{(2 \times (2+1))}{2} \right]^2 - 1 = 3^2 - 1 = 8 = 2^3 = \text{LHS}$ [1 mark]

Induction Step:

Assume $2^3 + 3^3 + \cdots + k^3 = \left[\frac{k(k+1)}{2} \right]^2 - 1$ for $k \geq 2$ [0.3 mark]

$$2^3 + 3^3 + \cdots + k^3 + (k+1)^3 = \left[\frac{k(k+1)}{2} \right]^2 - 1 + (k+1)^3 \quad [0.5 \text{ mark}]$$

$$= (k+1)^2 \left[\frac{k^2}{4} + k + 1 \right] - 1 = \frac{(k+1)^2}{4} [k^2 + 4k + 4] - 1 \quad [0.5 \text{ mark}]$$

$$= \frac{(k+1)^2}{2^2} \times (k+2)^2 - 1 = \left[\frac{(k+1)((k+1)+1)}{2} \right]^2 - 1 \quad [0.5 \text{ mark}]$$

Hence, by the Principle of Mathematical Induction, $2^3 + 3^3 + \cdots + n^3 = \left[\frac{n(n+1)}{2} \right]^2 - 1$ for $n \geq 2$.

..... [0.2 mark]

(d) Use the Principle of Mathematical Induction to prove that the inequality

$$1 + \frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{2^n} \geq 1 + \frac{n}{2}$$

for integers $n \geq 0$.

(3 marks)

Ans. Base Step:

When $n = 0$, LHS = $1 \geq 1 + \frac{0}{2} = \text{RHS}$ [1 mark]

[**Remark:** The left hand side is another notation of $\sum_{i=0}^{2^n} \frac{1}{i}$. Many students do not know how to prove this because **they refuse to put in any amount of efforts to understand series**]

Induction Step:

Assume $1 + \frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{2^k} \geq 1 + \frac{k}{2}$ for $n = k \geq 0$ [0.3 mark]

We now look at the series with 2^{k+1} terms

$$1 + \frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{2^k} + \underbrace{\frac{1}{2^k+1} + \cdots + \frac{1}{2^{k+1}}}_{2^{k+1} - 2^k = 2^k \text{ extra terms}}$$

$$\geq 1 + \frac{k}{2} + \underbrace{\frac{1}{2^k+1} + \cdots + \frac{1}{2^{k+1}}}_{2^{k+1} - 2^k = 2^k \text{ extra terms}} \quad [0.5 \text{ mark}]$$

$$\geq 1 + \frac{k}{2} + \left[\underbrace{\frac{1}{2^{k+1}} + \cdots + \frac{1}{2^{k+1}}}_{2^k \text{ extra terms}} \right] \left(\because \frac{1}{2^k+j} \geq \frac{1}{2^{k+1}} \text{ for } j = 1, \dots, 2^k \right) \quad [0.5 \text{ mark}]$$

$$= 1 + \frac{k}{2} + \frac{2^k}{2^{k+1}} = 1 + \frac{k}{2} + \frac{1}{2} = 1 + \frac{k+1}{2} \quad [0.5 \text{ mark}]$$

Hence, by the Principle of Mathematical Induction, $1 + \frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{2^n} \geq 1 + \frac{n}{2}$ for integers $n \geq 0$ [0.2 mark]

2. CO4. Express relations correctly with their mathematical properties. C2

(a) A set $A = \{1, 2, 3, 4\}$ has a relation

$$R = \{(2, 3), (3, 2), (4, 4)\}$$

- | | |
|--|----------------------------------|
| i. Is R irreflexive? | <input type="text" value="No"/> |
| ii. Is R symmetric? | <input type="text" value="Yes"/> |
| iii. Is R an equivalence relation? | <input type="text" value="No"/> |

Provide justifications for the above questions if your answer is No. ($3 \times 0.3 + 0.6 = 1.5$ mark)

Ans. i. R is not irreflexive because $(4, 4) \in R$ [0.3 mark]
 iii. R is not an equivalence relation because it is not reflexive, i.e. $(1, 1) \notin R$. [0.3 mark]

(b) Let A be all the factors of 60. The partial order \preceq for any $p, q \in A$ is defined by $p \preceq q$ if p is divisible by q . Sketch the Hasse diagram of the poset (A, \preceq) . (1.5 marks)

Ans. From $60 = 2^2 \times 3 \times 5$, we have


$$A = \{1, 2, 3, 5, 4, 6, 10, 15, 12, 20, 30, 60\}.$$

and the partial order is given by

$$\preceq = \{(p, q) : p \text{ is divisible by } q\}$$

For example, $(60, 2) \in \preceq$, i.e. 60 is divisible by 2.

The Hasse diagram is



List out all elements of A in proper ordering [0.7 mark]

Correct links in the Hasse diagram [0.8 mark]

(c) Consider a set $A = \{\alpha, \beta, \gamma, \delta, \epsilon, \zeta\}$ with the following relation

$$R = \{(\alpha, \delta), (\beta, \beta), (\gamma, \epsilon), (\delta, \zeta), (\epsilon, \gamma), (\zeta, \alpha)\}$$

- i. Write down the **matrix representation** of R and then (I) determine if the relation R is **reflexive** with justification; (II) determine if the relation R is **anti-symmetric** with justifications. (2 marks)

$$\text{Ans. } M_R = \begin{matrix} & \begin{matrix} \alpha & \beta & \gamma & \delta & \epsilon & \zeta \end{matrix} \\ \begin{matrix} \alpha \\ \beta \\ \gamma \\ \delta \\ \epsilon \\ \zeta \end{matrix} & \begin{pmatrix} 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \end{matrix} \dots\dots\dots [1 \text{ mark}]$$

Since $(\beta, \beta) \in R$, R is **not reflexive**. $\dots\dots\dots [0.5 \text{ mark}]$

Since $(\gamma, \epsilon), (\epsilon, \gamma) \in R$, R is **not anti-symmetric**. $\dots\dots\dots [0.5 \text{ mark}]$

- ii. Apply the **Warshall algorithm** to find the matrix representation of the **transitive closure** of R . (3 marks)

$$\text{Ans. Step 1: } M_R = \begin{matrix} & \begin{matrix} \alpha & \beta & \gamma & \delta & \epsilon & \zeta \end{matrix} \\ \begin{matrix} \alpha \\ \beta \\ \gamma \\ \delta \\ \epsilon \\ \zeta \end{matrix} & \begin{pmatrix} 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 1 & 0 & 0 \end{pmatrix} \end{matrix} \dots\dots\dots [0.5 \text{ mark}]$$

$$\text{Step 2: } M_R = \begin{matrix} & \begin{matrix} \alpha & \beta & \gamma & \delta & \epsilon & \zeta \end{matrix} \\ \begin{matrix} \alpha \\ \beta \\ \gamma \\ \delta \\ \epsilon \\ \zeta \end{matrix} & \begin{pmatrix} 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 1 & 0 & 0 \end{pmatrix} \end{matrix} \dots\dots\dots [0.5 \text{ mark}]$$

$$\text{Step 3: } M_R = \begin{matrix} & \begin{matrix} \alpha & \beta & \gamma & \delta & \epsilon & \zeta \end{matrix} \\ \begin{matrix} \alpha \\ \beta \\ \gamma \\ \delta \\ \epsilon \\ \zeta \end{matrix} & \begin{pmatrix} 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 1 & 0 & 0 \end{pmatrix} \end{matrix} \dots\dots\dots [0.5 \text{ mark}]$$

$$\text{Step 4: } M_R = \begin{array}{c} \alpha \quad \beta \quad \gamma \quad \delta \quad \epsilon \quad \zeta \\ \alpha \left(\begin{array}{cccccc} 0 & 0 & 0 & 1 & 0 & 1 \\ \beta \left(\begin{array}{cccccc} 0 & 1 & 0 & 0 & 0 & 0 \\ \gamma \left(\begin{array}{cccccc} 0 & 0 & 0 & 0 & 1 & 0 \\ \delta \left(\begin{array}{cccccc} 0 & 0 & 0 & 0 & 0 & 1 \\ \epsilon \left(\begin{array}{cccccc} 0 & 0 & 1 & 0 & 1 & 0 \\ \zeta \left(\begin{array}{cccccc} 1 & 0 & 0 & 1 & 0 & 1 \end{array} \right) \end{array} \right) \end{array} \right) \end{array} \right) \end{array} \dots [0.5 \text{ mark}]$$

$$\text{Step 5: } M_R = \begin{array}{c} \alpha \quad \beta \quad \gamma \quad \delta \quad \epsilon \quad \zeta \\ \alpha \left(\begin{array}{cccccc} 0 & 0 & 0 & 1 & 0 & 1 \\ \beta \left(\begin{array}{cccccc} 0 & 1 & 0 & 0 & 0 & 0 \\ \gamma \left(\begin{array}{cccccc} 0 & 0 & 1 & 0 & 1 & 0 \\ \delta \left(\begin{array}{cccccc} 0 & 0 & 0 & 0 & 0 & 1 \\ \epsilon \left(\begin{array}{cccccc} 0 & 0 & 1 & 0 & 1 & 0 \\ \zeta \left(\begin{array}{cccccc} 1 & 0 & 0 & 1 & 0 & 1 \end{array} \right) \end{array} \right) \end{array} \right) \end{array} \right) \end{array} \dots [0.5 \text{ mark}]$$

$$\text{Step 6: } M_R = \begin{array}{c} \alpha \quad \beta \quad \gamma \quad \delta \quad \epsilon \quad \zeta \\ \alpha \left(\begin{array}{cccccc} 1 & 0 & 0 & 1 & 0 & 1 \\ \beta \left(\begin{array}{cccccc} 0 & 1 & 0 & 0 & 0 & 0 \\ \gamma \left(\begin{array}{cccccc} 0 & 0 & 1 & 0 & 1 & 0 \\ \delta \left(\begin{array}{cccccc} 1 & 0 & 0 & 1 & 0 & 1 \\ \epsilon \left(\begin{array}{cccccc} 0 & 0 & 1 & 0 & 1 & 0 \\ \zeta \left(\begin{array}{cccccc} 1 & 0 & 0 & 1 & 0 & 1 \end{array} \right) \end{array} \right) \end{array} \right) \end{array} \right) \end{array} \dots [0.5 \text{ mark}]$$